

Separable Kernel of Nucleon-Nucleon Interaction in the Bethe-Salpeter Approach

Serge G. Bondarenko* and Valery V. Burov†

BLTP, Joint Institute for Nuclear Research, 141980 Dubna, Russia

Naohide Hamamoto‡ and Hiroshi Toki§

RCNP, Osaka University, Mihogaoka, Ibaraki, Osaka, 567-0047, Japan

The dispersion relations for nucleon-nucleon (NN) T -matrix in the framework of the Bethe-Salpeter equation for a two spin-one-half particle system and with a separable kernel of interaction are considered. The developed expressions are applied for construction of the separable kernel of interaction for S partial-waves in singlet and triplet channels. We calculate the low energy scattering parameters, the phase shifts, and the deuteron binding energy with the separable interaction. The approach can be easily extended to higher partial-waves for NN -scattering and other reactions ($\bar{N}N$ -, πN - scattering).

PACS numbers: 11.10.St, 13.75.Cs, 21.45.+v

I. INTRODUCTION

Recent and planned experiments on NN -scattering [1] as well as reactions with the deuteron and light nuclei reach large momentum transfer from the probe particle to the investigated system. In this context, relativistic effects play an essential role in understanding reaction mechanisms. Studies of elastic electron-deuteron scattering (see, for example, [2, 3]) in the nonrelativistic framework require to take into account mesonic exchange currents (a part of them has a pure relativistic origin, i.e. the so-called pair currents [4]). Theoretical calculations should therefore follow relativistic approaches.

The approach based on the Bethe-Salpeter equation (BS approach) [5] gives a powerful method to investigate various nuclear reactions both for electromagnetic and strong interactions. To find an analytic solution for the BS equation for T -matrix of NN scattering describing bound and scattering states, the separable kernel of interaction is very useful [6]. The separable kernel is well known from nonrelativistic consideration to solve the Lippman-Schwinger equation analytically [7]. The separable form of interaction is good not only for the technical reason but, in our opinion, also for the reflection of the nonlocal type of elementary nucleon-nucleon interaction (it was stressed first in papers [6]). This allows us not only to avoid the divergence peculiar of local theories but also to take into account the internal structure of the nucleon.

In the paper, we consider the BS equation for T -matrix and solve it using the separable kernel of interaction for a spin-one-half particle system (for a spin-zero case see, for example, [9] and for trinucleon systems [8]). This allows us to solve the BS equation without referring to the ladder approximation. We also find dispersion relations for the obtained solution which allow us to perform explicit analytic calculations and connect internal parameters of the kernel with experimental data: low-energy parameters, bound state energy, and phase shifts.

The paper is organized as follows: after the partial-wave decomposition of the BS equation (section II), the one-rank separable kernel of interaction is introduced and used for solving the BS equation (section III). Dispersion relations for the T -matrix are discussed in section IV. In section V, it is shown that the separable kernel of interaction includes the kernel in the ladder approximation, and the connection between parameters of two kernels is found. Section VI is devoted to results of calculation of the low-energy parameters, bound state energy, and phase shifts of elastic NN -scattering and to the discussion for the future works.

*Electronic address: bondarenko@jinr.ru

†Electronic address: burov@thsun1.jinr.ru

‡Electronic address: naohide@rcnp.osaka-u.ac.jp

§Electronic address: toki@rcnp.osaka-u.ac.jp

II. PARTIAL-WAVE DECOMPOSITION OF THE BS EQUATION

The BS equation for the two-nucleon T -matrix with momenta $p_1(p'_1), p_2(p'_2)$ could be represented in the following form:

$$T(p', p; P) = V(p', p; P) + \frac{i}{(2\pi)^4} \int dk V(p', k; P) S(k; P) T(k, p; P), \quad (1)$$

where $P = p_1 + p_2$, $p = (p_1 - p_2)/2$ [$p' = (p'_1 - p'_2)/2$] are total and relative momenta, and $V(p', p; P)$ is the NN kernel of interaction. The two-body propagator in terms of the total and relative momenta has the form:

$$S(k; P) = 1 / \left[(\hat{P}/2 + \hat{k} - m + i\epsilon)(\hat{P}/2 - \hat{k} - m + i\epsilon) \right], \quad (2)$$

with $\hat{a} = a_\mu \cdot \gamma^\mu$, and γ^μ are Dirac matrices.

After the partial wave decomposition in the rest frame of the two-nucleon system, the equation (1) becomes ($s = P^2$),

$$\begin{aligned} T_{\alpha\beta}(p'_0, p', p_0, p; s) &= V_{\alpha\beta}(p'_0, p', p_0, p; s) \\ &+ \frac{i}{4\pi^3} \sum_{\gamma\delta} \int dk_0 \int k^2 dk V_{\alpha\gamma}(p'_0, p', k_0, k; s) S_{\gamma\delta}(k_0, k; s) T_{\alpha\beta}(k_0, k, p_0, p; s). \end{aligned} \quad (3)$$

(Here and below $p \equiv |\mathbf{p}|$, $p' \equiv |\mathbf{p}'|$, $k \equiv |\mathbf{k}|$). Greek letters denote partial waves in the channels under consideration. In $LSJ\rho$ -basis [10, 11], the singlet 1S_0 channel consists of four partial waves, and the triplet 3S_1 channel, of eight partial-waves.

Below we omit all states which have at least one negative-energy nucleon and take only $^1S_0^+$ -wave in the singlet channel and neglect also $^3D_1^+$ -wave and take only $^3S_1^+$ -wave in the triplet channel. Therefore eq. (3) could be rewritten as:

$$\begin{aligned} t(p'_0, p', p_0, p; s) &= v(p'_0, p', p_0, p; s) \\ &+ \frac{i}{4\pi^3} \int dk_0 \int k^2 dk \frac{v(p'_0, p', k_0, k; s) t(k_0, k, p_0, p; s)}{(\sqrt{s}/2 - e_k + i\epsilon)^2 - k_0^2}, \end{aligned} \quad (4)$$

where $t \equiv T_{\tilde{\alpha}\tilde{\alpha}}$, $v \equiv V_{\tilde{\alpha}\tilde{\alpha}}$, with $\tilde{\alpha} = ^1S_0^+$ or $^3S_1^+$, and $e_k = \sqrt{k^2 + m^2}$ and m is the nucleon mass.

The T -matrix normalization condition could be written in the on-mass-shell form. Introducing phase shifts $\delta(s)$, one could write

$$t(s) \equiv t(0, \bar{p}, 0, \bar{p}, s) = -\frac{16\pi}{\sqrt{s}\sqrt{s - 4m^2}} e^{i\delta(s)} \sin \delta(s), \quad (5)$$

with $\bar{p} = \sqrt{s/4 - m^2} = \sqrt{2mT_{lab}}$.

To introduce also the low-energy parameters of NN -scattering, it is suitable to expand the above expression into a series of $\bar{p}$ -terms with the following expression [12]:

$$\bar{p} \cot \delta(s) = -a_0^{-1} + \frac{r_0}{2} \bar{p}^2 + \mathcal{O}(\bar{p}^3). \quad (6)$$

Taking into consideration only the first two terms of the decomposition (6), we define the scattering length a_0 and the effective radius r_0 .

As for the mass of the bound state (M_b), the squared T -matrix has a simple pole in the total momentum squared (s), and the bound state conditions could be written in the following form:

$$t(p'_0, p', p_0, p; s) = \frac{B(p'_0, p', p_0, p; s = M_b^2)}{s - M_b^2} + R(p'_0, p', p_0, p; s), \quad (7)$$

where functions B and R are regular at the point $s = M_b^2$. The bound state energy E_b is connected with M_b : $M_b = 2m - E_b$.

III. RANK I SEPARABLE KERNEL OF INTERACTION

The *ansatz* for the separable kernel of interaction with rank I has the form:

$$v(p'_0, p', p_0, p; s) = \lambda g(p'_0, p') g(p_0, p). \quad (8)$$

Here introduced is the g -function for NN -vertices, and λ is the coupling parameter.

To solve (4) we could assume the following separable form for T -matrix:

$$t(p'_0, p', p_0, p; s) = \tau(s) g(p'_0, p') g(p_0, p) \quad (9)$$

and substituting (8) and (9) into (4), we obtain the function $\tau(s)$:

$$\tau(s) = 1/(\lambda^{-1} + h(s)), \quad (10)$$

where the function $h(s)$ is:

$$h(s) = -\frac{i}{4\pi^3} \int dk_0 \int k^2 dk \frac{[g(k_0, k)]^2}{(\sqrt{s}/2 - e_k + i\epsilon)^2 - k_0^2}. \quad (11)$$

Therefore, the T -matrix could be rewritten in the following form:

$$t(p'_0, p', p_0, p; s) = \frac{g(p'_0, p') g(p_0, p)}{\lambda^{-1} + h(s)}, \quad (12)$$

and the on-mass-shell expression is:

$$t(s) = \frac{n(s)}{d(s)} = \frac{[g(0, \bar{p})]^2}{\lambda^{-1} + h(s)}. \quad (13)$$

It should be noted that the expression (13) has the so-called N/D -form widely used in the nonrelativistic T -matrix theory [12] and in some relativistic methods.

Using (13), it is easy to connect the solution of T -matrix and the phase shifts $\delta(s)$. To achieve this, we assume that the imaginary part of the function $n(s)$ satisfies the following condition:

$$\text{Im } n(s) = 0. \quad (14)$$

This condition is connected with the specific choice of g -functions for NN -vertices to be discussed in the next section. By using (13) and (14), the phase shifts $\delta(s)$ could be expressed by the formula:

$$\cot \delta(s) = \frac{\text{Re } t(s)}{\text{Im } t(s)} = -\frac{\lambda^{-1} + \text{Re } h(s)}{\text{Im } h(s)} \quad (15)$$

To express the low-energy parameters in terms of the T -matrix solution, it is suitable to expand the function $h(s)$ in a series of $\bar{p}$ terms:

$$h(s) = h_0 + i\bar{p}h_1 + \bar{p}^2h_2 + i\bar{p}^3h_3 + \mathcal{O}(\bar{p}^4), \quad (16)$$

$$\text{Re } h(s) = h_0 + \bar{p}^2h_2 + \mathcal{O}(\bar{p}^4), \quad (17)$$

$$\text{Im } h(s) = \bar{p}(h_1 + \bar{p}^2h_3 + \mathcal{O}(\bar{p}^3)). \quad (18)$$

Using now the definition (6) and (15-18), we obtain the a_0 and r_0 parameters:

$$a_0 = \frac{h_1}{\lambda^{-1} + h_0}, \quad r_0 = \frac{2}{h_1} \left[(\lambda^{-1} + h_0) \frac{h_3}{h_1} - h_2 \right]. \quad (19)$$

The bound state condition (7) with the help of (12) is now expressed in the form:

$$\lambda^{-1} = -h(s = M_b^2). \quad (20)$$

IV. DISPERSION RELATIONS

Let us consider now the analytic properties of the BS solution (namely, the function h). The simplest choice of the function $g(p_0, p)$ is the *Yamaguchi* type [6]:

$$g(p_0, p) = (p_0^2 - p^2 - \beta^2 + i\epsilon)^{-1}. \quad (21)$$

In this case, the function h could be rewritten as

$$h(s, \beta) = -\frac{i}{4\pi^3} \partial_{\beta^2} \int dp_0 \int p^2 dp \frac{1}{(\sqrt{s}/2 - e_p + i\epsilon)^2 - p_0^2} \frac{1}{p_0^2 - e_\beta^2 + i\epsilon} \quad (22)$$

with $\partial_{\beta^2} \equiv \partial/\partial\beta^2$ and $e_\beta = \sqrt{p^2 + \beta^2}$. We have introduced the second argument β to extract the explicit dependence of the function h on this parameter.

Analyzing equation (22), we could conclude that there are four poles in the complex plane p_0 , namely:

$$\begin{aligned} p_0^{(1)}(s) &= \frac{\sqrt{s}}{2} - e_p + i\epsilon & p_0^{(2)}(s) &= -\frac{\sqrt{s}}{2} + e_p - i\epsilon \\ p_0^{(3)}(s) &= -e_\beta + i\epsilon & p_0^{(4)}(s) &= e_\beta - i\epsilon \end{aligned} \quad (23)$$

With the change of the variable s , the poles $p_0^{(1)}$ and $p_0^{(3)}$ move, and there could be a situation when two poles “pinch” real p_0 axes. It means, the function $h(s)$ has at this s -point the leap and imaginary part. First points at which this condition is satisfied (branch points) could be found from the following equations:

$$\begin{aligned} p_0^{(1)}(s) &= -p_0^{(1)}(s) & \Rightarrow & & s_0 &= 4m^2, \\ p_0^{(1)}(s) &= -p_0^{(2)}(s) & \Rightarrow & & s_1 &= 4(m + \beta)^2. \end{aligned} \quad (24)$$

Summarizing the situation, one could say that the function $h(s)$ has two cuts starting at points s_0 and s_1 , respectively, and therefore could be written in a dispersion form:

$$\begin{aligned} h(s, \beta) &= \int_{4m^2}^{+\infty} \frac{\rho(s', \beta) ds'}{s' - s - i\epsilon}, \\ \rho(s', \beta) &= \theta(s' - 4m^2) \rho_{el}(s', \beta) + \theta(s' - 4(m + \beta)^2) \rho_{in}(s', \beta) \end{aligned} \quad (25)$$

with two spectral functions $\rho_{el;in}$ (*el* stands for *elastic* and *in* for *inelastic*) which are connected with the imaginary parts as

$$\rho(s', \beta) = \frac{1}{\pi} \text{Im } h(s', \beta) = \frac{1}{2\pi i} (h - h^*). \quad (26)$$

To find spectral functions, one should perform p_0 -integration in (22) which gives the following result:

$$h(s, \beta) = -\frac{1}{2\pi^2} \partial_{\beta^2} \int p^2 dp \frac{1}{s/4 - \sqrt{s}e_p + m^2 - \beta^2 + i\epsilon} \left[\frac{1}{\sqrt{s} - 2e_p + i\epsilon} + \frac{1}{2e_\beta} \right]. \quad (27)$$

Taking into account the following symbolic equation:

$$\frac{1}{x - x_0 \pm i\epsilon} = \frac{\mathcal{P}}{x - x_0} \mp i\pi\delta(x - x_0), \quad (28)$$

it is easy to find the spectral functions:

$$s' \geq 4m^2 \quad (29)$$

$$\rho_{el}(s', \beta) = \sqrt{s'} \sqrt{s' - 4m^2} / (\pi^2 (s' - 4m^2 + 4\beta^2)^2) \quad (30)$$

$$s' \geq 4(m + \beta)^2 \quad (31)$$

$$\begin{aligned} \rho_{in}(s', \beta) &= -(64\beta^6 + 16\beta^4 s' - 192\beta^4 m^2 - 20\beta^2 s'^2 \\ &\quad + 192\beta^2 m^4 - 32\beta^2 s' m^2 + 16m^4 s' + 3s'^3 - 64m^6 - 12m^2 s'^2) / \\ &\quad (2\pi^2 s' (s' - 4m^2 + 4\beta^2)^2 \sqrt{s' - 4(m + \beta)^2} \sqrt{s' - 4(m - \beta)^2}). \end{aligned}$$

To perform integration (22), it is suitable to introduce new variables:

$$\lambda_\beta = \frac{\beta}{m}, \quad t = \frac{s}{4m^2} \quad (32)$$

$$\rho^2 = 1 - \frac{4m^2}{s} = 1 - \frac{1}{t}, \quad w^2 = \frac{m^2}{m^2 - \beta^2} = \frac{\lambda_\beta^2}{1 - \lambda_\beta^2}, \quad (33)$$

$$v^2 = \frac{m + \beta}{m - \beta} = \frac{1 + \lambda_\beta}{1 - \lambda_\beta}, \quad \sigma^2 = -\frac{s - 4(m + \beta)^2}{s - 4(m - \beta)^2} = -\frac{t - (1 + \lambda_\beta)^2}{t - (1 - \lambda_\beta)^2}. \quad (34)$$

In case when the following conditions are valid

$$m > \beta > 0 \quad \Rightarrow \quad 1 > \lambda_\beta > 0, \quad (35)$$

$$4(m + \beta)^2 > s > 4m^2 \quad \Rightarrow \quad (1 + \lambda_\beta)^2 > t > 1, \quad (36)$$

the introduced parameters are real and positive:

$$\begin{aligned} 1 > \rho^2 > 0, & \quad w^2 > 0, \\ v^2 > 1 > 0, & \quad \sigma^2 > 0. \end{aligned} \quad (37)$$

The condition (36) means that the second (inelastic) imaginary part does not contribute to the function $\text{Im } h(s)$ when the phase shifts are calculated in the region $4(m + \beta)^2 > s > 4m^2$, and therefore:

$$\text{Im } h(s, \beta) = \text{Im } h_{el}(s, \beta) = \frac{\rho(\rho^2 - 1)(1 + w^2)^2}{4m^2\pi(\rho^2 + w^2)^2}, \quad \text{if } 4(m + \beta)^2 > s > 4m^2. \quad (38)$$

Performing integration (25), we obtain for the real part of function $h(s)$ (the imaginary part is given by (38)):

$$\begin{aligned} \text{Re } h(s, \beta) &= h_{el}(s, \beta) + h_{in}(s, \beta), \\ h_{el}(s, \beta) &= \frac{\rho(\rho^2 - 1)(1 + w^2)^2}{4m^2\pi^2(\rho^2 + w^2)^2} \ln \left| \frac{\rho + 1}{\rho - 1} \right| \\ &\quad - \frac{(\rho^2 - 1)(1 + w^2)^2(w^2 - \rho^2)}{4m^2\pi^2(\rho^2 + w^2)^2 w} \arctan 1/w \\ &\quad - \frac{(\rho^2 - 1)(1 + w^2)}{4m^2\pi^2(\rho^2 + w^2)}, \\ h_{in}(s) &= -\frac{(1 + \sigma^2)(1 + v^2)^2}{32m^2\pi^2(v^4 + \sigma^2)} \ln \left| \frac{v^2 + 1}{v^2 - 1} \right| \\ &\quad - \frac{(1 + \sigma^2)(1 + v^2)^2}{16m^2\pi^2(v^2 - 1)^2 v(v^2 - \sigma^2)^2} \\ &\quad \times (v^6 + v^4\sigma^2 - 4v^4 + 4v^2\sigma^2 - v^2 - \sigma^2) \arctan 1/v \\ &\quad + \frac{(1 + \sigma^2)(1 + v^2)^3}{16m^2\pi^2\sigma(v^4 + \sigma^2)(v^2 - \sigma^2)^2\pi^2(v^2 - 1)^2} \\ &\quad \times (-v^6 + 2v^6\sigma^2 - 3v^4\sigma^2 + 3\sigma^4v^2 - 2\sigma^4 + \sigma^6) \arctan 1/\sigma \\ &\quad - \frac{(1 + \sigma^2)(1 + v^2)^2}{16m^2\pi^2(v^2 - 1)(v^2 - \sigma^2)} \end{aligned} \quad (39)$$

To find also expressions for low-energy parameters, we should return to (18)-(19) and expand the function $h(s)$ in a series of $\bar{p}$ terms:

$$\begin{aligned} h_0(\beta) &= \frac{(1 + w^2)}{4m^2\pi^2w^3} (w + (1 + w^2) \arctan 1/w) \\ &\quad + \frac{5 + 20v^2 + 5v^8 + 14v^4 + 20v^6}{4m^2\pi^2(v^2 - 1)^3(3v^2 + 1)} \sqrt{\frac{3v^2 + 1}{v^2 + 3}} \arctan 1/\sqrt{\frac{3v^2 + 1}{v^2 + 3}} \\ &\quad - \frac{(v^2 + 1)^4}{4m^2\pi^3v(v^2 - 1)^3} \arctan 1/v - \frac{1}{8m^2\pi^2} \ln \left| \frac{v^2 + 1}{v^2 - 1} \right| - \frac{(v^2 + 1)^2}{4m^2\pi^2(v^2 - 1)^2} \end{aligned} \quad (40)$$

$$h_2(\beta) = -\frac{(1+w^2)^2}{4m^4\pi^2w^5}(3w+(3+w^2)\arctan 1/w) \quad (41)$$

$$\begin{aligned} & -\frac{1}{4m^4\pi^2(v^2-1)^5(v^2+3)(3v^2+1)^2}(304v^2+1212v^{12}+3790v^8 \\ & +304v^{14}+2704v^{10}+29v^{16}+1212v^4+2704v^6+29) \\ & \times \sqrt{\frac{3v^2+1}{v^2+3}} \arctan 1 / \sqrt{\frac{3v^2+1}{v^2+3}} \\ & + \frac{(v^4+10v^2+1)(v^2+1)^4}{4m^4\pi^2v(v^2-1)^5} \arctan 1/v \\ & + \frac{1}{8m^4\pi^2} \ln \left| \frac{v^2+1}{v^2-1} \right| + \frac{11v^8+52v^6+66v^4+52v^2+11)(v^2+1)^2}{8m^4\pi^2(v^2-1)^4(3v^2+1)(v^2+3)} \\ h_1(\beta) &= \frac{(1+w^2)^2}{4m^3\pi w^4} \quad (42) \end{aligned}$$

$$h_3(\beta) = -\frac{(1+w^2)^2(4+3w^2)}{8m^5\pi w^6} \quad (43)$$

At this point we have *explicit analytic* expressions which connect the parameters of the separable kernel of interaction (λ and β [λ_β]) with observables: phase shifts $\delta(s)$ (eq. (15)), low-energy parameters a_0 and r_0 (eq. (19)), and bound state energy (eq. (20)).

V. SEPARABLE AND ONE-MESON EXCHANGE KERNELS OF INTERACTION

In this section, we show that the so-called realistic kernel of interaction (meson-nucleon) in the ladder approximation is included in the separable kernel of the Yamaguchi-type g-functions. To achieve this, we introduce a simple approximation for the ladder kernel:

$$V(p_0, p; k_0, k) \rightarrow \tilde{V}(p_0, p; k_0, k) = \frac{V(p_0, p; 0, 0)V(0, 0; k_0, k)}{V(0, 0; 0, 0)}. \quad (44)$$

Using the expressions of the kernel for scalar-meson (sc) exchange [10],[13]:

$$V_{sc}(p_0, p; k_0, k) = -\frac{g_{sc}^2}{4\pi} \frac{1}{\pi^2} \frac{1}{4pk e_p e_k} [(e_p e_k + m^2)Q_0(z_\mu) - pk Q_1(z_\mu)], \quad (45)$$

where $z_\mu = (p^2 + k^2 - (p_0 - k_0)^2 + \mu^2)/(2pk)$, μ is the exchange meson mass, and $Q_i(z)$ are the Legendre functions of second kind, $Q_0(z) = 1/2 \ln(z+1)/(z-1)$, $Q_1(z) = zQ_0(z) - 1$, it is clear that

$$V_{sc}(p_0, p; 0, 0) = \lim_{k_0 \rightarrow 0, k \rightarrow 0} V_{sc}(p_0, p; k_0, k) = a_p g_\mu(p_0, p), \quad (46)$$

with

$$\begin{aligned} a_p &= \frac{g_{sc}^2}{4\pi} \frac{1}{\pi^2} \frac{e_p + m}{2e_p}, \\ g_\mu(p_0, p) &= 1/(p_0^2 - p^2 - \mu^2). \end{aligned} \quad (47)$$

Expressions for $V_{sc}(0, 0; k_0, k)$ could be obtained from eqs. (46-47) by the following substitutions: $p_0 \rightarrow k_0$ and $p \rightarrow k$. To make connection between parameters more clear, we perform the p/m -decomposition in function a_p up to $\mathcal{O}(p^2/m^2)$ term:

$$\begin{aligned} a_p &= a_k = \frac{g_{sc}^2}{4\pi} \frac{1}{\pi^2}, \\ V_{sc}(0, 0; 0, 0) &= -\frac{g_{sc}^2}{4\pi} \frac{1}{\pi^2} \frac{1}{\mu^2}. \end{aligned} \quad (48)$$

Using now expression (44), we could write:

$$\tilde{V}_{sc}(p_0, p; k_0, k) = -\frac{g_{sc}^2}{4\pi} \left(\frac{\mu}{\pi}\right)^2 g_\mu(p_0, p) g_\mu(k_0, k). \quad (49)$$

Comparing the above expressions with the separable form of the kernel introduced by (8), we could write the following connection between parameters:

$$\beta = \mu, \quad \lambda_{sc} = -\frac{g_{sc}^2}{4\pi} \left(\frac{\mu}{\pi}\right)^2. \quad (50)$$

Equations (50) are valid also for the vector-meson (vc) exchange kernel with the substitution of $g_{sc} \rightarrow g_{vc}$ and μ by the vector-meson mass and could be derived from the following expression:

$$V_{vc}(p_0, p; k_0, k) = -\frac{g_{vc}^2}{4\pi} \frac{1}{\pi^2} \frac{1}{4pk e_p e_k} [-2(2e_p e_k + m^2)Q_0(z_\mu)] . \quad (51)$$

The pseudoscalar-meson (ps) exchange kernel of interaction has the form:

$$V_{ps}(p_0, p; k_0, k) = -\frac{g_{sc}^2}{4\pi} \frac{1}{\pi^2} \frac{1}{4pk e_p e_k} [-(e_p e_k - m^2)Q_0(z_\mu) + pk Q_1(z_\mu)] , \quad (52)$$

and for $p, k \rightarrow 0$ we could write:

$$\tilde{V}_{ps}(p_0, p; k_0, k) \sim p^2 k^2 g_\mu(p_0, p) g_\mu(k_0, k). \quad (53)$$

In this case, the expression for $\tilde{V}_{ps}$ goes to zero for $p, k \rightarrow 0$, and it is impossible to find a connection between parameters similar to (50).

We have obtained connections between parameters for various cases of the scalar and vector exchange, which are given in Ref. [13]. Results are given in Table I.

VI. RESULTS AND DISCUSSION

At this moment, we could find internal parameters of the kernel of interaction (λ, β) to reproduce experimental values for low-energy parameters $a_{0s}^{exp}, r_{0s}^{exp}$ for the singlet channel (1S_0) and a_{0t}^{exp} and bound state (deuteron) energy E_d^{exp} for the triplet channel (3S_1).

In the case of 1S_0 -channel, we use equation (19) with $a_0 \equiv a_{0s}^{exp}$ to find λ :

$$\lambda^{-1} = (a_{0s}^{exp})^{-1} h_1(\beta) - h_0(\beta). \quad (54)$$

Inserting the obtained expression into equation (20) with $r_0 \equiv r_{0s}^{exp}$, we find:

$$r_{0s}^{exp} = \frac{2}{h_1(\beta)} [(a_{0s}^{exp})^{-1} h_3(\beta) - h_2(\beta)] . \quad (55)$$

Solving the nonlinear equation (55), we could find the value of β and, then, using expression (54), the value of λ .

In the case of 3S_1 -channel, we obtain λ from the bound state condition (20) with $E_b \equiv E_d^{exp}$:

$$\lambda^{-1} = -h(s = (M_d^{exp})^2, \beta), \quad (56)$$

where $M_d = 2m - E_d$. Inserting the obtained expression into (19) with $a_0 \equiv a_{0t}^{exp}$, we find:

$$a_{0t}^{exp} = \frac{h_1(\beta)}{h_0(\beta) - h(s = M_d^2, \beta)}. \quad (57)$$

Solving the nonlinear eq. (57), we could find the value of β and, then, using expression (56) - the value of λ .

To solve equations (55) and (57), we use FORTRAN code with the DZEROX subroutine (as a part of CERNLIB package). As a result, we find the following values for λ and β :

$$\begin{aligned} &\text{for } ^1S_0 \text{ channel :} \\ &\quad \lambda = -0.29425404 \text{ GeV}^{-2}, \quad \beta = 0.22412880 \text{ GeV}, \\ &\text{for } ^3S_1 \text{ channel :} \\ &\quad \lambda = -0.79271213 \text{ GeV}^{-2}, \quad \beta = 0.27160579 \text{ GeV}, \end{aligned} \quad (58)$$

TABLE I: Connection between parameters of two kernels.

J^P		μ (GeV)	$g^2/4\pi$	λ (GeV ²)
0^+	$NN\delta$	0.983	0.64	-.06265954891
0^+	$NN\sigma'$	0.550	7.07	-.2166930823
1^-	$NN\rho$	0.769	0.43	-.02576448048
1^-	$NN\omega$	0.7826	10.6	-.6577877886

TABLE II: The binding energy and low-energy parameters for singlet and triplet channels.

$1S_0$	a_{0s} (Fm)	r_{0s} (Fm)	
Calculated	-23.748	2.75	
Experiment	-23.748 ± 0.010	2.75 ± 0.05	
$3S_1$	a_{0t} (Fm)	r_{0t} (Fm)	E_d (MeV)
Calculated	5.424	1.774	2.224644
Experiment	5.424 ± 0.004	1.759 ± 0.005	2.224644 ± 0.000046

The results on the binding energy of the deuteron and the low-energy parameters are shown in Table II. The experimental data are taken from [14].

The phase shifts $\delta_s(s)$ and $\delta_t(s)$ calculated with the above parameters are presented in Fig. 1. The experimental data are taken from [15]. As it is seen from the figure, even with the simplest choice of the separable kernel of interaction - rank I with only two parameters λ and β , we find the low-energy parameters of elastic NN scattering a_s and r_s in the singlet channel and a_t and bound state (deuteron) energy E_d in the triplet channel with required accuracy and reproduce phase shifts in a region till $T_{lab} = 100 - 150$ MeV.

To summarize, we recall once again the basic points of our lines of thought and the approximations made in the analysis. We start with the Bethe-Salpeter equation for T -matrix and perform the partial-wave decomposition. At this point, we omit all states which have at least one negative-energy nucleon and consider only $^1S_0^+$ -wave in the singlet channel and neglect also $^3D_1^+$ -wave and left only $^3S_1^+$ -wave in the triplet channel. In order to find the explicit analytic solution for T -matrix, we introduce separable ansatz for the kernel of interaction. As a result, we find the N/D -form for on-shell T -matrix. Taking the simplest choice of g -function in a *Yamaguchi* form for NN -vertex, we analyze the analytic structure of the obtained solution for T -matrix and write the dispersion representation of the T -matrix. Performing the integration in the dispersion formula, we obtain *explicit analytic* expressions which connect the parameters of the separable kernel of interaction (λ and β [λ_β]) with observables: phase shifts, low-energy parameters and bound state energy. This connection is used to calculate intrinsic parameters of the kernel of interaction and then to find phase shifts.

We find also that the kernel of interaction in the ladder approximation is included into the separable kernel. To obtain this result, we use a simple approximation for the expressions in the one-meson exchange model.

The main result of the paper is the dispersion form of T -matrix for elastic NN scattering obtained for the separable kernel of interaction which allows us to perform analytic calculations and to connect explicitly parameters of the kernel and observables. The discussed method could be technically expanded to the multi-rank separable kernels and more complex forms of g -function for NN -vertex to achieve required accuracy for bound state energy, low-energy parameters and phase shifts in a wide region of cm laboratory energy. The approach can be easily extended to higher partial-waves for NN -scattering and other reactions ($\bar{N}N$ -, πN - scattering).

Acknowledgments. We would like to thank Prof. Y. Yamaguchi and Prof. Y. Nambu for useful discussions. One of us (V.V.B.) expresses his deep gratitude to the directorate of RCNP for hospitality. The work was supported in part by the RFBR grant N 00-15-96737.

REFERENCES

- [1] C. Lechanoine-LeLuc, F. Lehar, Rev. of Modern Phys., **65** (1993), p. 47.
- [2] Gari M., Hyuga H., Z.Phys. A **58** (1976), p.291.
- [3] V.V. Burov, V.N. Dostovalov, and S.Eh.Sus'kov, Czech. J. of Phys. **41** (1991) p. 1139; Particles and Nuclei **23** (1992), p.721.




FIG. 1: The singlet δ_s and triplet δ_t phase shifts.

- [4] S.G. Bondarenko, V.V. Burov, M. Beyer, S.M. Dorkin, Phys. Rev. C **58** (1998) 3143-3152.
- [5] E.E. Salpeter and H.A. Bethe, Phys. Rev. C **84** (1951) p.1232.
- [6] Y. Yamaguchi, Phys. Rev. **95**, (1954) p.1628; Y. Yamaguchi, Y. Yamaguchi, Phys. Rev. **95**, (1954) p.1635.
- [7] L. Mathelitsch, W. Plessas, W. Schweiger, Phys. Rev. C **26** (1982) p.65; W. Schweiger, J. Haidenbauer, W. Plessas, Phys. Rev. C **32** (1985), p.1261.
- [8] G. Rupp and J.A. Tjon, Phys. Rev. C **37**, 1729 (1988); G. Rupp and J.A. Tjon, Phys. Rev. C **41**, 472 (1990); G. Rupp, Nucl. Phys. A **508**, 131c (1990); G. Rupp and J.A. Tjon, Phys. Rev. C **45**, 2133 (1992); G. Rupp, Proc. of PAN XIII, 13th Int. Conf. on Particles and Nuclei, Perugia, June-July 1993, World Scientific, Ed. A. Pascolini, p. 673-675 (1994).
- [9] K. Schwarz *et al.*, Acta Physica Austriaca, **53** (1981), p.191; J. Fröhlich *et al.*, Phys. Rev. C **25** (1982), p.2591; J. Fröhlich *et al.*, Phys. Rev. C **27** (1983), p.265; J. Haidenbauer, and J. Fröhlich, Phys. Rev. C **33**, 456 (1986).
- [10] J.J. Kubis, Phys. Rev. D **6** (1972) p.547.
- [11] S.G. Bondarenko, V.V. Burov, M. Beyer, S.M. Dorkin, Few Body Syst. **26** (2-4) (1999) p.185.
- [12] G.E. Brown, A.D. Jackson, "The Nucleon-nucleon interaction", North-Holland Publishing Company, 1976.
- [13] R. Machleidt *et al.*, Phys. Rep. **149** (1987) p.1.
- [14] O. Dumbrajs *et al.*, Nucl. Phys. B **216** (1983) p.277.
- [15] R. A. Arndt, L. D. Roper, R. A. Bryan, R. B. Clark, B. J. VerWest, and P. Signell, Phys. Rev. D **28** (1983) p.97; R.A. Arndt, J.S. Hyslop and L.D. Roper, Phys. Rev. D **35** (1987) p.128.